

On the Existence of Smooth Solutions to the Dyachenko Equations Governing Free-Surface Unsteady Ideal Fluid Flows

R. V. Shamin

Presented by Academician V.E. Zakharov June 3, 2005

Received June 6, 2005

DOI: 10.1134/S1064562406010303

Let an inviscid incompressible fluid occupy a domain in the plane (x, y) bounded by the free surface

$$\begin{aligned} -\infty < y \leq \eta(x, t), \\ -\infty < x < \infty, \quad t > 0. \end{aligned}$$

Assuming that the fluid flow is potential, we have $\mathbf{v}(x, y, t) = \nabla\Phi(x, y, t)$, where $\mathbf{v}(x, y, t)$ is a two-dimensional velocity field and $\Phi(x, y, t)$ is a potential. Throughout this paper, the gradient operator and Laplacian are applied with respect to the spatial variables x and y only. The incompressibility condition $\operatorname{div} \mathbf{v} = 0$ implies that the velocity potential obeys the Poisson equation

$$\Delta\Phi(x, y, t) = 0. \quad (1)$$

Equation (1) is supplemented with the boundary and initial conditions

$$(\eta_t + \Phi_x \eta_x - \Phi_y)|_{y=\eta(x,t)} = 0, \quad (2)$$

$$\left(\Phi_t + \frac{1}{2} |\nabla\Phi|^2 + gy \right) \Big|_{y=\eta(x,t)} = 0, \quad (3)$$

$$\Phi_y|_{y=-\infty} = 0, \quad (4)$$

$$\eta|_{t=0} = \eta_0(x), \quad (5)$$

$$\Phi|_{t=0} = \Phi_0(x, y), \quad (6)$$

where g is the acceleration of gravity.

The first results on the existence of analytical solutions to these problems were obtained in [1]. The existence of solutions of finite smoothness was proved in [2, 3].

Numerical methods for these problems were discussed in [4–6]. However, the original equations written in terms of (η, Φ) are not very convenient in the numerical modeling of free surface problems.

In [7, 8] conformal transformations were used to study free surface problems. Systems of integro-differential equations solved for the time derivatives were obtained. Equivalent equations, called the Dyachenko equations, were derived in [8]. These equations were found to be very convenient for numerical solution.

In this paper, we prove the existence of analytical solutions to the Dyachenko equations on a sufficiently small time interval. It is shown also that these solutions on the real line belong to Sobolev spaces, which is important for substantiating numerical methods.

For positive s , let $Q_s = \{w = u + iv \in \mathbb{C} : 0 < u < 2\pi, -\infty < v < s\}$ be an unbounded domain and E_s^0 denote the space of square integrable analytic functions in Q_s that are 2π -periodic with respect to u . The space E_s^0 equipped with the inner product

$$(f, g)_{E_s^0} = \int_{Q_s} f(w) \overline{g(w)} dw$$

is a Hilbert space (see [9, ch. 1, Theorem 1]).

Let $f(w) \in E_s^0$. Then $f(w) = \sum_{k=1}^{\infty} f_k e^{-ikw}$, where the

series converges in the E_s^0 norm. Since Q_s is unbounded, the functions equal to a constant do not belong to E_s^0 . We introduce the space $E_s = E_s^0 + \mathbb{R}$

equipped with the norm $\|f\|_{E_s} = \left(\sum_{k=0}^{\infty} |f_k|^2 \right)^{1/2}$.

Shirshov Institute of Oceanology, Russian Academy of Sciences, Nakhimovskii pr. 36, Moscow, 117851 Russia
e-mail: roman@shamin.ru

The operator $P: E_s \times E_s \rightarrow E_s$ is defined as follows.

Let $A = \sum_{k=0}^{\infty} a_k e^{-ikw}$ and $B = \sum_{k=0}^{\infty} b_k e^{-ikw}$. Then

$$P(A, B) = \frac{c_0}{2} + \sum_{k=1}^{\infty} c_k e^{kv} e^{-iku},$$

where $c_k = \sum_{m=k}^{\infty} a_m \bar{b}_{m-k}$.

Consider the Dyachenko equations

$$\begin{aligned} R_t &= i(UR' - UR), \quad w \in Q_s, \quad t \in (0, T), \\ V_t &= i(UV' - B'R) + g(R - 1), \end{aligned} \tag{7}$$

where $U = P(R, V) + P(V, R)$, $B = P(V, V)$, and the prime denotes the derivative with respect to the complex variable w . Equations (7) are subject to the boundary conditions

$$\begin{aligned} R, V &\text{ are } 2\pi\text{-periodic with respect to } u, \\ |R| &\rightarrow 1, \quad |V| \rightarrow 0, \quad u \rightarrow -\infty, \quad v < 0 \end{aligned} \tag{8}$$

and the initial conditions

$$R(w, 0) = R_0(w), \quad V(w, 0) = V_0(w), \quad w \in Q_s. \tag{9}$$

The initial functions are assumed to belong to E_s .

Definition 1. Functions R and V that are analytic with respect to t and take their values in E_s are called an analytical solution to problem (7)–(9) if these functions satisfy (7)–(9).

Theorem 1. Let $R_0, V_0 \in E_{s_1}$, where $s_1 > 0$. Then, for any $0 < s < s_1$, there is $T(s) > 0$ such that problem (7)–(9) has a unique analytical solution $R, V \in E_s$ for $t < T(s)$.

The proof of the theorem is based on the Nirenberg–Nishida theorem (see [10, p. 220 in the Russian edition]).

In numerical simulation, it is convenient to deal with solutions on the real line in Sobolev spaces.

The relevant function spaces are defined as follows. Let $\Omega = (0, 2\pi) \times (0, T)$ with $0 < T < \infty$ be a rectangle in the plane (u, t) . By $W_2^k(\Omega)$, we denote the Sobolev

space of complex-valued functions from $L_2(\Omega)$ with all generalized derivatives up to the k th order from $L_2(\Omega)$, equipped with the norm

$$\|u\|_{W_2^k(\Omega)} = \left\{ \sum_{|\alpha| \leq k} \int_{\Omega} |D^\alpha u(x)|^2 dx \right\}^{\frac{1}{2}}.$$

Theorem 2. Let R, V be an analytical solution for $t < T$.

Then, at $v = 0$ $R, V \in W_2^k(\Omega)$ for any $k \geq 1$.

ACKNOWLEDGMENTS

The author is grateful to Academician V.E. Zakharov and A.I. Dyachenko for suggesting this problem and their interest in this work.

This work was supported by the Russian Foundation for Basic Research (project nos. 04-05-64784, 04-01-00256) and the program for basic research of the Pre-sidium of the Russian Academy of Sciences “Mathematical Methods in Nonlinear Dynamics.”

REFERENCES

1. V. I. Nalimov, Dokl. Akad. Nauk SSSR **189**, 45–49 (1969).
2. V. I. Nalimov, Sib. Mat. Zh. **37**, 1356–1366 (1996).
3. V. I. Nalimov, Din. Splosh. Sredy **18**, 104–210 (1974).
4. V. V. Voinov and O. V. Voinov, Dokl. Akad. Nauk SSSR **221**, 559–562 (1975).
5. A. I. Petrov and V. G. Smolyanin, Prikl. Mat. Mekh. **57** (1), 137–143 (1993).
6. R. M. Garipov, Dokl. Akad. Nauk SSSR **161**, 547–550 (1965).
7. A. I. Dyachenko, V. E. Zakharov, and E. A. Kuznetsov, Fiz. Plazmy **22**, 916–928 (1996) [Plasma Phys. Rep. **22**, 829–840 (1996)].
8. V. E. Zakharov, A. I. Dyachenko, and O. A. Vasilyev, Eur. J. Mech. **21**, 283–291 (2002).
9. D. Gaier, *Vorlesungen über approximation im komplexen* (Birkhäuser, Basel, 1980; Mir, Moscow, 1986).
10. L. Nirenberg, *Topics in Nonlinear Functional Analysis* (Courant Institute, New York Univ., New York, 1974; Mir, Moscow, 1977).